

ON THE HOMOTOPY INVARIANCE OF L^2 TORSION FOR COVERING SPACES.

VARGHESE MATHAI AND MELVIN ROTHENBERG

ABSTRACT. We prove the homotopy invariance of L^2 torsion for covering spaces, whenever the covering transformation group is either residually finite or amenable. In the case when the covering transformation group is residually finite and when the L^2 cohomology of the covering space vanishes, the homotopy invariance was established by Lück [Lu]. We also give some applications of our results.

INTRODUCTION

L^2 analytic torsion for L^2 acyclic covering spaces with positive decay was first studied in [M] and in [L] and L^2 Reidemeister-Franz torsion was first studied in [CM], see also [LuR], [Lu]. These L^2 torsion invariants were subsequently generalised by using the theory of determinant lines of finitely generated Hilbertian modules over finite von Neumann algebras, which was initiated in [CFM]. Non-zero elements of the determinant lines can be viewed as volume forms on the Hilbertian modules. Using this, they constructed both L^2 combinatorial torsion and L^2 analytic torsion invariants associated to flat Hilbertian bundles of determinant class over compact polyhedra and manifolds, as volume forms on the L^2 homology and L^2 cohomology respectively, under the assumption that the covering space was of determinant class. These L^2 torsion invariants specialise to the Ray-Singer-Quillen torsion and the classical Reidemeister-Franz torsion respectively, in the finite dimensional case. Using the results of [BFKM], it was shown in [CFM] that the combinatorial and analytic L^2 torsions were equal whenever the covering space is of determinant class. By this result and calculations done in [M] and [L], one obtains a simplicial description of hyperbolic volume for a closed 3-dimensional hyperbolic manifold in terms of the L^2 Reidemeister-Franz torsion.

In this paper, we prove the homotopy invariance of L^2 torsion for covering spaces, whenever the covering transformation group is either residually finite or amenable. In the case when the covering transformation group is residually finite and when the L^2 cohomology of the covering space vanishes, the homotopy invariance was established by Lück [Lu]. Using our main theorem, we can define the L^2 torsion of a discrete group Γ , where Γ is either residually finite or amenable and such that $B\Gamma$ is a finite CW complex. The L^2 torsion of the discrete group Γ is then an element of the determinant line of the reduced L^2 cohomology of the group Γ . We also give a new proof of the classical result that the hyperbolic volume of a closed

Date: MARCH 1997.

1991 Mathematics Subject Classification. Primary: 58G11, 58G18 and 58G25.

Key words and phrases. L^2 torsion, invariants, amenable groups, residually finite groups, Whitehead groups, homotopy invariance.

Mel Rothenberg was supported in part by NSF Grant DMS 9423300.

3-dimensional hyperbolic manifold is a homotopy invariant, which actually only uses the part of our theorem which was proved by Lück [Lu]. Our paper relies on the recent results that residually finite covering spaces are of determinant class due to [Lu], [BFK] and that amenable covering spaces are of determinant class due to [DM2].

The authors thank Shmuel Weinberger for some helpful conversations.

1. PRELIMINARIES

1.1. Determinants and determinant lines. Here we will review some of the results on the Fuglede-Kadison determinant and on determinant lines of Hilbertian modules over finite von Neumann algebras. The results of this section are mainly from the paper [CFM]. See also the papers [CM], [FK], [Lu1] and [LuR].

Let $\mathcal{A}$ be a finite von Neumann algebra with a fixed faithful finite normal trace $\tau : \mathcal{A} \rightarrow \mathbb{C}$. Let k denote the involution on $\mathcal{A}$. Then $\mathcal{A}$ has a scalar product $\langle a, b \rangle = \tau(b * a)$ for $a, b \in \mathcal{A}$. Let $\ell^2(\mathcal{A})$ denote the completion of $\mathcal{A}$ with respect to this scalar product.

A *Hilbert $\mathcal{A}$ module* is a Hilbert space M together with a continuous left $\mathcal{A}$ module structure such that there is an isometric $\mathcal{A}$ -linear embedding of M into $\ell^2(\mathcal{A}) \otimes H$ for some Hilbert space H ($\mathcal{A}$ acts trivially on H). Note that the embedding is *not* part of the structure. M is said to be *finitely generated* if H can be chosen above to be finite dimensional.

A Hilbert module has a particular scalar product. If we ignore the scalar product, we get what we call a Hilbertian $\mathcal{A}$ module, more precisely a *Hilbertian $\mathcal{A}$ module* is a topological vector space M with a left $\mathcal{A}$ -module structure such that there is a scalar product $\langle, \rangle$ on M which generates the topology of M , and such that $(M, \langle, \rangle)$ is a Hilbert $\mathcal{A}$ module.

Any scalar product $\langle, \rangle$ on M as above will be called *admissible*. Suppose that $\langle, \rangle$ is another scalar product on M . Then there is an operator $A : M \rightarrow M$ such that $\langle v, w \rangle_1 = \langle Av, w \rangle$ which satisfies

- (1) A is a linear homeomorphism since $\langle, \rangle$ and $\langle, \rangle_1$ define the same topology,
- (2) self adjoint with respect to $\langle, \rangle$,
- (3) positive,
- (4) A commutes with $\mathcal{A}$.

Any two admissible scalar products on a Hilbertian $\mathcal{A}$ -module give rise to isomorphic Hilbert $\mathcal{A}$ -modules. In particular, if we choose an admissible scalar product on M , it becomes a Hilbert $\mathcal{A}$ module and the *von Neumann dimension* $\dim_\tau(M)$, which is defined as the von Neumann trace (defined in the next paragraph) of the orthogonal projection from $\ell^2(\mathcal{A}) \otimes H$ onto M , is independent of the choice of admissible scalar product.

The *commutant* of M , $B(M)$ is the algebra of all bounded linear operators commuting with $\mathcal{A}$. Then there is a canonical trace on $B(M)$, Tr_τ defined as follows: If M were free, is isomorphic to $\ell^2(\mathcal{A}) \otimes \mathbb{C}^k$, then $B(M)$ consists of $k \times k$ matrices with entries in $\mathcal{A}$, acting on the right and for $f \in B(M)$,

$$\text{Tr}_\tau(f) = \sum_{i=1}^k \tau(f_{ii})$$

is the trace. In the general case, $\text{Tr}_\tau(f) = \text{Tr}_\tau(i_M \circ f \circ \pi_M)$, where i_M denotes the embedding of M into a free module and π_M denotes the projection from the free module onto M .

Let $GL(M)$ denote the group of all *invertible operators* in the commutant of M , $B(M)$. Then there is a determinant function,

$$\text{Det}_\tau : GL(M) \rightarrow \mathbb{R}^+$$

called the *Fuglede-Kadison determinant*, which satisfies

- (1) $\text{Det}_\tau(AB) = \text{Det}_\tau(A)\text{Det}_\tau(B)$
- (2) $\text{Det}_\tau(\lambda I) = |\lambda|^{\dim_\tau(M)}$ for $\lambda \in \mathbb{C}$
- (3) $\text{Det}_{\lambda\tau}(A) = \text{Det}_\tau(A)^\lambda$ for $\lambda > 0$
- (4) Det_τ is continuous in the norm topology on $GL(M)$

Its definition is as follows [CFM]: Let $A \in GL(M)$ and A_t be a piecewise smooth path from the identity to A (since $GL(M)$ is connected, cf. [Di]). Define

$$(*) \quad \log \text{Det}_\tau(A) = \int_0^1 \Re \text{Tr}_\tau(A_t^{-1} \dot{A}_t) dt.$$

where $\Re$ denotes the real part. Using a result of Araki, Smith and Smith [ASS], if $t \rightarrow A_t$ is a loop, then $\int_0^1 \Re \text{Tr}_\tau(A_t^{-1} \dot{A}_t) dt = 0$, so that $(*)$ is well defined.

Let M be a finitely generated Hilbertian $\mathcal{A}$ -module. Define the *determinant line* of M , denoted by $\det M$, to be the real vector space generated by the symbols $\langle, \rangle$, one for each *admissible* scalar product on M , with the following relations : $\langle, \rangle_1$, $\langle, \rangle_2$ are admissible scalar products on M , then

$$\langle, \rangle_2 = \text{Det}_\tau(A)^{-1/2} \langle, \rangle_1$$

where $A \in GL(M)$, $A > 0$ and

$$\langle v, w \rangle_2 = \langle Av, w \rangle_1 \quad \forall v, w \in M.$$

It has been shown in [CFM] that $\det(M)$ is a real *one dimensional* vector space.

1.2. L^2 torsion for covering spaces. We will discuss L^2 torsion in the special case of normal covering spaces; for the general case of Hilbertian $(\mathcal{A} - \pi)$ bimodules, we refer to [CFM], [BFKM].

Let $\Gamma \rightarrow \widehat{M} \rightarrow M$ be a normal covering space of a compact manifold. Let K be a smooth triangulation of M , and $\widehat{K}$ be the induced Γ -invariant triangulation of $\widehat{M}$. Then there is a natural inner product on the space of finite support co-chains on $\widehat{K}$, $C^j(\widehat{K})$ where we declare the simplices to be an orthonormal basis. Then Γ acts by isometries on $C^j(\widehat{K})$ and we consider the Hilbert space completion $C_{(2)}^j(\widehat{K})$. The coboundary operator

$$d^K : C^j(\widehat{K}) \rightarrow C^{j+1}(\widehat{K})$$

induces a bounded operator

$$d^{\widehat{K}} : C_{(2)}^j(\widehat{K}) \rightarrow C_{(2)}^{j+1}(\widehat{K}).$$

Let $d^{\widehat{K}*}$ denote the L^2 adjoint of $d^{\widehat{K}}$, and the combinatorial Laplacian

$$\Delta_j^{\widehat{K}} = d^{\widehat{K}} d^{\widehat{K}*} + d^{\widehat{K}*} d^{\widehat{K}} : C_{(2)}^j(\widehat{K}) \rightarrow C_{(2)}^j(\widehat{K})$$

is a bounded self adjoint operator, which commutes with the Γ -action, since $d^{\widehat{K}}$ and $d^{\widehat{K}*}$ do. Let $\Delta_j^{\widehat{K}+}$ denote the restriction of $\Delta_j^{\widehat{K}}$ onto the orthogonal complement of

$\ker \Delta_j^{\widehat{K}}$. Assume now that $\widehat{M}$ is of *determinant class*, that is, the Fuglede-Kadison determinant of $\Delta_j^{\widehat{K}+}$, $\text{Det}_\tau(\Delta_j^{\widehat{K}+})$ is defined and is strictly positive for all $j \geq 0$, [CM], [BFKM]. It has been shown that $\widehat{M}$ is of determinant class whenever Γ is either residually finite (cf. [Lu], [BFK]) or amenable (cf. [DM2]).

Since $\ker \Delta_j^{\widehat{K}} \subset C_{(2)}^j(\widehat{K})$, it inherits an admissible scalar product. By the combinatorial analogue of the Hodge theorem, one sees that $\ker \Delta_j^{\widehat{K}}$ is isomorphic to the reduced L^2 cohomology $\bar{H}_{(2)}^j(\widehat{K})$, it follows that $\bar{H}_{(2)}^j(\widehat{K})$ gets an admissible scalar product, i.e. an element in the determinant line $\det(\bar{H}_{(2)}^j(\widehat{K}))$. The alternating tensor product of these elements is an element in $\det(\bar{H}_{(2)}^\bullet(\widehat{K}))$, which we denote by

$$\hat{\varphi} = \hat{\varphi}(K) \in \det(\bar{H}_{(2)}^\bullet(\widehat{K})).$$

The L^2 Reidemeister-Franz torsion, $\varphi_{\widehat{K}} \in \det(\bar{H}_{(2)}^\bullet(\widehat{K}))$ is defined as in [CFM] to be

$$\varphi_{\widehat{K}} = \prod_{j=0}^n \text{Det}_\tau(\Delta_j^{\widehat{K}+})^{\frac{(-1)^j j}{2}} \hat{\varphi}(K).$$

Theorem 1.1 (Topological invariance of L^2 Reidemeister-Franz torsion [CFM]). *Assume that $\widehat{M}$ be of determinant class. Then $\varphi_{\widehat{K}}$ is a topological invariant of M , whenever $\dim M$ is odd.*

Remark 1.2. When $\bar{H}_{(2)}^j(\widehat{K}) = 0 \ \forall j \geq 0$, this result was first obtained by [CM]. In fact, $\varphi_{\widehat{K}}$ is an invariant of the simple homotopy type of K , see [LuR] and 2.1 below. In the even dimensional case, the L^2 Reidemeister-Franz torsion is not a topological invariant in general. Also, if $\dim M$ is even and $\bar{H}_{(2)}^j(\widehat{K}) = 0 \ \forall j \geq 0$, then $\varphi_{\widehat{K}} = 1$.

There is also an analytic version of L^2 Reidemeister-Franz torsion which can be defined in an analogous manner. It is called L^2 *analytic torsion* and denoted by $\varphi_{\widehat{M}} \in \det(\bar{H}_{(2)}^\bullet(\widehat{M}))$. We refer to [CFM] for details of its definition.

Theorem 1.3 (Metric invariance of L^2 analytic torsion [CFM]). *Assume that $\widehat{M}$ be of determinant class. Then $\varphi_{\widehat{M}}$ is independent of the choice of Riemannian metric g on M whenever $\dim M$ is odd.*

Remark 1.4. When $\bar{H}_{(2)}^j(\widehat{M}) = 0 \ \forall j \geq 0$, this result was first obtained by [M] and [L]. In [CFM], an even more general result is obtained. In the even dimensional case, the L^2 analytic torsion does in general depend on the choice of Riemannian metric. Also, if $\dim M$ is even and $\bar{H}_{(2)}^j(\widehat{M}) = 0 \ \forall j \geq 0$, then $\varphi_{\widehat{M}} = 1$.

Under the de Rham isomorphism, which identifies $\bar{H}_{(2)}^j(\widehat{K})$ and $\bar{H}_{(2)}^j(\widehat{M})$ for all j (see [Do]), the results of [BFKM] are used to prove the following theorem in [CFM].

Theorem 1.5 (Equality of L^2 torsions [CFM], [BFKM]). *Assume that $\widehat{M}$ be of determinant class. Then via the identification of determinant lines induced by the de Rham isomorphism, the L^2 analytic torsion and L^2 Reidemeister-Franz torsion are equal, that is, $\varphi_{\widehat{M}} = \varphi_{\widehat{K}}$, whenever the dimension of M is odd.*

2. HOMOTOPY INVARIANCE

Here we establish the homotopy invariance of L^2 torsion for residually finite covering transformation groups and for amenable covering transformation groups. As mentioned before, the case when the L^2 cohomology vanishes and when the covering transformation group is residually finite, this result is due to [Lu].

We first formulate the problem of the homotopy invariance of L^2 torsion as follows. Let $f : M \rightarrow N$ be a homotopy equivalence of compact manifolds, $\widehat{f} : \widehat{M} \rightarrow \widehat{N}$ be the induced homotopy equivalence of the normal Γ -covering spaces $\widehat{M}$ and $\widehat{N}$. Then $\widehat{f}$ induces an isomorphism on the reduced L^2 cohomology, $\widehat{f}^* : \bar{H}_{(2)}^\bullet(\widehat{N}) \rightarrow \bar{H}_{(2)}^\bullet(\widehat{M})$. By 2.3 in [CFM], $\widehat{f}^*$ induces an isomorphism $\widehat{f}_*$ on determinant lines, $\widehat{f}_* : \det \bar{H}_{(2)}^\bullet(\widehat{N}) \rightarrow \det \bar{H}_{(2)}^\bullet(\widehat{M})$. Thus a homotopy equivalence of manifolds induces a canonical isomorphism of determinant lines of L^2 cohomology.

Conjecture 1 (Homotopy invariance of L^2 torsion). *Let $f : M \rightarrow N$ be a homotopy equivalence of compact odd dimensional manifolds. Suppose that $\widehat{M}$ is of determinant class (or equivalently if $\widehat{N}$ is of determinant class). Via the identification of determinant lines of L^2 cohomology of normal Γ covering spaces as above, we conjecture that*

$$\phi_{\widehat{M}} = \phi_{\widehat{N}} \in \det \bar{H}_{(2)}^\bullet(\widehat{M})$$

This conjecture was first formulated by Lück [Lu] in the case when the L^2 cohomology of M is trivial.

Our first result will be a general formula, which generalises the ones in [LuR], [Lu] and [Lu1], in the case when the L^2 cohomology is trivial. Let $A \in GL(n, \mathbb{Z}[\Gamma])$ be an invertible matrix with entries in $\mathbb{Z}[\Gamma]$. Then it defines a bounded invertible operator $R_A : \ell^2(\Gamma)^n \rightarrow \ell^2(\Gamma)^n$ which commutes with the left action of Γ on $\ell^2(\Gamma)^n$. So it has a positive Fuglede-Kadison determinant $\text{Det}(R_A)$. Also A defines an element in the Whitehead group of Γ , and it can be shown that the Fuglede-Kadison determinant Det induces a homomorphism

$$\Phi_\Gamma : Wh(\Gamma) \rightarrow \mathbb{R}^+$$

which was defined in [LuR] and [Lu]. Now a homotopy equivalence $f : M \rightarrow N$ defines an acyclic complex of cochains over the group ring of Γ , $C^\bullet(M_f)$, where we now choose cell decomposition for M and N , and f to be a cellular homotopy equivalence and M_f is the cellular mapping cone. Then as in [Mi], this acyclic complex defines the Whitehead torsion $T(f) \in Wh(\Gamma)$ of the homotopy equivalence f . Note that via the identification above, one has $\phi_{\widehat{M}} \otimes \phi_{\widehat{N}}^{-1} \in \mathbb{R}$. Then we have the following

Proposition 2.1. *Let $f : M \rightarrow N$ be a homotopy equivalence of compact odd dimensional manifolds. Suppose that $\widehat{M}$ is of determinant class. Via the identification of determinant lines of L^2 cohomology of normal Γ -covering spaces as above, one has*

$$\phi_{\widehat{M}} \otimes \phi_{\widehat{N}}^{-1} = \Phi_\Gamma(T(f)) \in \mathbb{R}^+$$

Remark 2.2. Farrell and Jones [FJ] have proved that the Whitehead group $Wh(\Gamma)$ is trivial whenever Γ is the fundamental group of a manifold with non-positive sectional curvature. It follows from Proposition 2.1 that in this case, one has

$$\phi_{\widehat{M}} = \phi_{\widehat{N}} \in \det \bar{H}_{(2)}^\bullet(\widehat{M})$$

i.e., the homotopy invariance of L^2 torsion, provided the covering spaces are of determinant class. Also Farrell and Jones [FJ] conjectured that the Whitehead group of any torsion-free group vanishes, which would imply Conjecture 2 on the homotopy invariance of L^2 torsion.

To prove this proposition, we recall some definitions and facts from [CFM] about abstract determinant class complexes. We begin with;

Definition 2.3. Consider a finitely generated Hilbertian module M over $\mathcal{A}$. A scalar product $\langle \cdot, \cdot \rangle$ on M will be called *D-admissible* if it can be represented in the form

$$\langle v, w \rangle = \langle A(v), w \rangle_1 \quad \text{for } v, w \in M,$$

where $\langle \cdot, \cdot \rangle_1$ is an admissible scalar product on M and $A \in B(M)$ is an injective (possibly not invertible) homomorphism $A : M \rightarrow M$, which is positive and self-adjoint with respect to $\langle \cdot, \cdot \rangle_1$, and the following property is satisfied: if

$$A = \int_0^\infty \lambda dE_\lambda$$

is the spectral decomposition of A and if

$$\phi(\lambda) = \dim_\tau(E_\lambda) = \text{Tr}_\tau(E_\lambda)$$

denotes the corresponding spectral density function, then the integral

$$\int_0^\infty \ln(\lambda) d\phi(\lambda) > -\infty$$

is assumed to converge to a finite value.

By a morphism of Hilbertian modules, we always mean one which preserves the modules structure over the von Neumann algebra. An injective homomorphism of Hilbertian modules $f : M \rightarrow N$ with dense image will be called a *D-isomorphism* if for some, and therefore all, admissible scalar product $\langle \cdot, \cdot \rangle_N$ on N , the induced scalar product $\langle \cdot, \cdot \rangle_M$ on M given by $\langle v, m \rangle_M = \langle f(v), f(m) \rangle_N$, is *D-admissible*. Note that any *D-isomorphism* of Hilbertian modules $f : M \rightarrow N$, induces an isomorphism of determinant lines $f_* : \det(M) \rightarrow \det(N)$, defined above, cf. [CFM].

A sequence of Hilbertian modules and morphisms $0 \rightarrow M' \xrightarrow{\alpha} M \xrightarrow{\beta} M'' \rightarrow 0$ is called *D-exact* if α is a monomorphism, $\text{im } \alpha = \ker \beta$ and the map induced by β , $M/\ker \beta \rightarrow M''$ is a *D-isomorphism*.

Let

$$0 \rightarrow C^0 \xrightarrow{d} C^1 \xrightarrow{d} \dots \xrightarrow{d} C^N \rightarrow 0$$

be a cochain complex of finite length formed by finitely generated Hilbertian $\mathcal{A}$ modules and bounded linear maps commuting with the action of $\mathcal{A}$. Here $\mathcal{A}$ is a finite von Neumann algebra, such as the group von Neumann algebra of Γ . Let Z^i denote the submodule of cocycles and B^i the submodule of coboundaries. Then the cochain complex $C^\bullet$ is said to be of *determinant class* if the following sequence

$$0 \rightarrow Z^i \xrightarrow{\alpha_i} C^i \xrightarrow{\beta_i} \bar{B}^i \rightarrow 0$$

is *D-exact*, that is $\text{im}(\alpha_i) = \ker(\beta_i)$ and the map induced by β_i , $C^i/\ker(\beta_i) \rightarrow \bar{B}^i$ is a *D-isomorphism*. This agrees with the notion of determinant class discussed in section 1.2, (cf. [CFM]).

We note as in [CFM] that the property that a cochain complex $C^\bullet$ of finitely generated Hilbertian modules to be of determinant class depends only on the homotopy type of $C^\bullet$ in the category of finitely generated complexes of Hilbertian modules over $\mathcal{A}$.

Definition 2.4. Let $C^\bullet$ be a cochain complex which is of determinant class. Then by Proposition 3.10 [CFM] there is a canonical isomorphism between the determinant lines of the graded Hilbertian modules,

$$\phi_C : \det(C) \rightarrow \det(\bar{H}(C))$$

where $\bar{H}(C)$ denotes the reduced L^2 cohomology. This gives a definition of the L^2 torsion of an abstract cochain complex $C^\bullet$ which is of determinant class,

$$\phi_C \in \det(C)^{-1} \otimes \det(\bar{H}(C))$$

Let K be a smooth triangulation on M , and $\widehat{K}$ be the induced Γ -invariant triangulation on $\widehat{M}$. It can be shown that the L^2 torsion of the L^2 cochain complex $C_{(2)}^\bullet(\widehat{K})$ of $\widehat{K}$ is equal to the L^2 Reidemeister-Franz torsion of $\widehat{K}$, (see Proposition 3.11 in [CFM]).

Proposition 2.5. *Let $f : C \rightarrow C'$ be a cochain homotopy equivalence of determinant class complexes. Then*

$$\phi_{C'}^{-1} \otimes \phi_C = \phi_{C_f} \in \det(C_f)^{-1}.$$

where C_f denotes the mapping cone complex.

Proof. There is a short exact sequence

$$0 \rightarrow C'^{-1} \rightarrow C_f \rightarrow C \rightarrow 0$$

where C_f denotes the mapping cone complex, and C'^{-1} denotes the complex C' shifted in degree by -1 . By Proposition 3.5 [CFM], one has the canonical isomorphisms

$$\det(C'^{-1}) \otimes \det(C) \rightarrow \det(C_f)$$

and

$$\det(\bar{H}(C'^{-1})) \otimes \det(\bar{H}(C)) \rightarrow \mathbb{C}$$

since C_f is acyclic. Therefore we get the canonical isomorphism

$$\det(C') \otimes \det(\bar{H}(C'))^{-1} \otimes \det(C)^{-1} \otimes \det(\bar{H}(C)) \rightarrow \det(C_f)^{-1}$$

which yields the identity

$$\phi_{C'}^{-1} \otimes \phi_C = \phi_{C_f} \in \det(C_f)^{-1}.$$

□

Proof of Proposition 2.1. By Proposition 2.5, it suffices to determine the quantity $\phi_{C_f} \in \det(C_f)^{-1}$ in our particular case. Firstly, since the complexes and the exact sequences are based, we see that $\phi_{C_f} \in \mathbb{R}$. Next, C_f defines the Whitehead torsion $T(f) \in Wh(\Gamma)$, and it is clear that the L^2 torsion ϕ_{C_f} is just the Fuglede-Kadison determinant of the Whitehead torsion $T(f)$, $\Phi_\Gamma(T(f))$. □

Our next result is for amenable groups.

Proposition 2.6. *Suppose that Γ is a finitely presented amenable group. Then the homomorphism*

$$\Phi_\Gamma : Wh(\Gamma) \rightarrow \mathbb{R}^+$$

is trivial.

Proof. Let $a \in Wh(\Gamma)$. Then by [Mi], it is represented as the Whitehead torsion of a homotopy equivalence $f : L \rightarrow K$ of finite CW complexes, which we can assume without loss of generality is an inclusion, i.e. $a = T(f)$. Let $\hat{K}$ and $\hat{L}$ denote the corresponding Γ normal covering complexes. The relative cochain complex $C(\hat{K}, \hat{L})$ is acyclic, and so is its L^2 completion $C_{(2)}(\hat{K}, \hat{L})$. In particular, the combinatorial Laplacian $\Delta_j^{\hat{K}, \hat{L}}$ is invertible, and we see that

$$\Phi_\Gamma(T(f)) = \prod_{j=0}^n \text{Det}_\tau(\Delta_j^{\hat{K}, \hat{L}})^{\frac{(-1)^j j}{2}} > 0.$$

Let $\mathcal{F}_K$ denote a fundamental domain for the action of the group Γ on $\hat{K}$, and let $\mathcal{F}_L = \mathcal{F}_K \cap \hat{L}$, which is a fundamental domain for the action of Γ on $\hat{L}$.

Since Γ is amenable, by the Følner criterion for amenability one gets a *regular exhaustion*, that is, a sequence $\{X_m\}_{m=1}^\infty$ of *finite* subsets of Γ such that

$$\lim_{m \rightarrow \infty} \frac{\#\partial_\delta X_m}{\#X_m} = 0$$

where $\partial_\delta X_m = \{\gamma \in \Gamma : d(\gamma, X_m) < \delta \text{ and } d(\gamma, \Gamma - X_m) < \delta\}$ is a δ -neighbourhood of the "boundary" of X_m , $d(\cdot, \cdot)$ denotes the word metric on Γ and $\#X_m$ denotes the number of elements in X_m .

One then gets *regular exhaustions* $\{L_m\}_{m=1}^\infty$ of $\hat{L}$, and $\{K_m\}_{m=1}^\infty$ of $\hat{K}$, that is a sequences of finite subcomplexes of $\hat{L}$ and $\hat{K}$ respectively such that

$$(1) \quad L_m = \bigcup_{g \in X_m} g \cdot \mathcal{F}_L \text{ and } K_m = \bigcup_{g \in X_m} g \cdot \mathcal{F}_K;$$

$$(2) \quad \hat{L} = \bigcup_{m=1}^\infty L_m \text{ and } \hat{K} = \bigcup_{m=1}^\infty K_m$$

(3) Let $\dot{N}_{m,\delta}$ denote the number of elements $g \in \Gamma$ which have distance (with respect to the word metric in Γ) less than or equal to δ from any element $g' \in \Gamma$ such that the intersection of $g' \cdot \mathcal{F}_L$ with the topological boundary ∂L_m of L_m is not empty, that is, $\dot{N}_{m,\delta} = \#\partial_\delta X_m$. Let $N_m = \#X_m$, which also equals the number of translates of $\mathcal{F}_L$ which are contained in L_m . Then by the amenability assumption, one has for every $\delta > 0$,

$$\lim_{m \rightarrow \infty} \frac{\dot{N}_{m,\delta}}{N_m} = 0.$$

Note that $\dot{N}_{m,\delta}$ also equals the number of elements $g \in \Gamma$ which have distance (with respect to the word metric in Γ) less than or equal to δ from any element $g' \in \Gamma$ such that the intersection of $g' \cdot \mathcal{F}_K$ with the topological boundary ∂K_m of K_m is not empty, and N_m also equals the number of translates of $\mathcal{F}_K$ which are contained in K_m .

We can then define the relative cochain complexes $C(K_m, L_m)$ (note that $L_m = K_m \cap \hat{L}$). Let $\Delta_j^{(m)}$ denote the combinatorial Laplacian acting on $C^j(K_m, L_m)$. Let $\Delta_j^{(m)+}$ denote the combinatorial Laplacian acting on the orthogonal complement of

its null-space. Then by the results of section 4 in [DM2], which are easily generalised to the relative situation which is being considered here, one has

$$\text{Det}_\tau(\Delta_j^{\hat{K}, \hat{L}}) \geq \limsup_{m \rightarrow \infty} \det_{\mathbb{C}}(\Delta_j^{(m)+})^{\frac{1}{2N_m}} = 1$$

since $\Delta_j^{(m)+}$ is an invertible matrix with integer entries. To prove equality, we notice that if $E_m(\lambda)$ denotes the spectral density function of $\Delta_j^{(m)}$, then

$$(1) \quad F_m(\lambda) = F_m(0) \quad \text{for all } \lambda < K^{-2}, m \geq 0.$$

where $F_m(\lambda) = \frac{E_m(\lambda)}{N_m}$ and K is a constant as in Lemma 2.2 in [DM2] such that

$$K^2 \geq \max\{\|\Delta_j^{\hat{K}, \hat{L}}\|, \|\Delta_j^{(m)+}\|, \|\Delta_j^{(m)+^{-1}}\|, \|\Delta_j^{\hat{K}, \hat{L}^{-1}}\|\}$$

for all $m \geq 0$. Actually in Lemma 2.2 in [DM2], one obtains a constant K_1 such that

$$K_1^2 \geq \max\{\|\Delta_j^{\hat{K}, \hat{L}}\|, \|\Delta_j^{(m)+}\|\}$$

for all $m \geq 0$. Then an identical proof as in Lemma 2.2 in [DM2] gives the existence of a constant K_2 such that

$$K_2^2 \geq \max\{\|\Delta_j^{(m)+^{-1}}\|, \|\Delta_j^{\hat{K}, \hat{L}^{-1}}\|\}.$$

Finally, choose K to be the greater of K_1, K_2 .

The estimate (1) says that $\Delta_j^{(m)}$ has a spectral gap at zero which is bounded away from zero, independent of $m \geq 0$. Let $\bar{F}(\lambda) = \limsup_{m \rightarrow \infty} F_m(\lambda)$, where we assume without loss of generality that $\bar{F}(\lambda)$ is right continuous. By Theorem 2.1 in [DM2], one has $\bar{F}(\lambda) = F(\lambda)$, where $F(\lambda)$ denotes the spectral density function of $\Delta_j^{\hat{K}, \hat{L}}$. Therefore

$$\begin{aligned} 0 \leq \bar{F}(\lambda) - \bar{F}(0) &= \limsup_{m \rightarrow \infty} F_m(\lambda) - \lim_{m \rightarrow \infty} F_m(0) \\ &\leq \sup\{|F_m(\lambda) - F_m(0)| : m \geq 0\} = 0 \end{aligned}$$

if $\lambda < K^{-2}$, by (1). Therefore one has,

$$(2) \quad F(\lambda) = \bar{F}(\lambda) = \bar{F}(0) \quad \text{for } \lambda < K^{-2}$$

The estimate (2) says that $\Delta_j^{\hat{K}, \hat{L}}$ also has a spectral gap of size at least K^{-2} , at zero. Also observe that

$$\int_0^{K^2} \sup\{F_m(\lambda) - F_m(0) : m \geq 0\} d\lambda = \int_{K^{-2}}^{K^2} \sup\{F_m(\lambda) - F_m(0) : m \geq 0\} d\lambda < \infty.$$

By the dominated convergence theorem, one sees as in Lemma 3.3.1 in [Lu] that

$$\int_0^{K^2} \frac{F(\lambda) - F(0)}{\lambda} d\lambda = \lim_{m \rightarrow \infty} \int_0^{K^2} \frac{F_m(\lambda) - F_m(0)}{\lambda} d\lambda.$$

It follows that

$$\text{Det}_\tau(\Delta_j^{\hat{K}, \hat{L}}) = \lim_{m \rightarrow \infty} \det_{\mathbb{C}}(\Delta_j^{(m)+})^{\frac{1}{2N_m}} = 1.$$

This completes the proof of the Proposition. $\square$

Assembling Propositions 2.1 and 2.6 in this section, and using Theorem 0.5, part a) in [Lu], which states that

$$\Phi_\Gamma : Wh(\Gamma) \rightarrow \mathbb{R}^+$$

is trivial whenever Γ is residually finite, and using Theorem 1.7 of the previous section, we obtain the following theorem which gives significant evidence to the conjecture stated at the beginning of the section.

Theorem 2.7. *Let $f : M \rightarrow N$ be a homotopy equivalence of compact, odd dimensional manifolds. Via the identification of determinant lines of L^2 cohomology of normal Γ covering spaces as in the beginning of the section, and whenever Γ is either residually finite or Γ is amenable, one has*

$$\phi_{\widehat{M}} = \phi_{\widehat{N}} \in \det \bar{H}_{(2)}^\bullet(\widehat{M}).$$

3. APPLICATIONS

In this section, we briefly give some applications of Theorem 2.7.

3.1. L^2 torsion for discrete groups. Our first application is the definition of the L^2 torsion for a certain class of discrete groups.

Proposition 3.1. *Let Γ be either a residually finite group or an amenable group, whose classifying space $B\Gamma$ is a finite CW complex of odd dimension. Then one can define the L^2 torsion of Γ as*

$$\phi_\Gamma = \phi_{E\Gamma} \in \det(\bar{H}_{(2)}^\bullet(\Gamma)).$$

Proof. We note that since Γ be either a residually finite group or an amenable group such that $B\Gamma$ is a finite CW complex, it follows that $E\Gamma$ is of determinant class, and therefore the L^2 torsion $\phi_{E\Gamma}$ is well defined. Now $B\Gamma$ is only well defined upto homotopy, and since both the determinant class condition depends only on the homotopy type of $B\Gamma$, and by Theorem 2.7, the L^2 torsion is a homotopy invariant, we therefore see that the L^2 torsion $\phi_{E\Gamma}$ is independent of the choice of $B\Gamma$, and depends only on the group Γ . $\square$

We now conjecture the following.

Conjecture 2. *Let Γ be a discrete group whose classifying space $B\Gamma$, is a finite CW complex of odd dimension. Then one can define the L^2 torsion of Γ as*

$$\phi_\Gamma = \phi_{E\Gamma} \in \det(\bar{H}_{(2)}^\bullet(\Gamma)).$$

3.2. Homotopy invariance of hyperbolic volume. Our next application is a new proof of the homotopy invariance of the hyperbolic volume of closed 3-dimensional hyperbolic manifolds. Although it uses Theorem 2.7, it uses only the part that was proved by Lück [Lu].

Proposition 3.2. *Let M and N be a homotopy equivalent closed 3-dimensional hyperbolic manifolds. Then their hyperbolic volumes are equal.*

Proof. By the calculations in [M] and [L], one sees that the reduced L^2 cohomology of M and N vanish, and that the universal covers $\widehat{M} = \widehat{N} = \mathbb{H}^3$, which is hyperbolic 3-space, is of determinant class, and finally that the L^2 torsions of $\widehat{M}$ and $\widehat{N}$ can be calculated in terms of the hyperbolic volume,

$$-6\pi \log \phi_{\widehat{M}} = \text{vol}(M)$$

and

$$-6\pi \log \phi_{\tilde{N}} = \text{vol}(N).$$

Now applying Theorem 1.5 and Theorem 2.7, we see that

$$\text{vol}(M) = -6\pi \log \phi_{\tilde{M}} = -6\pi \log \phi_{\tilde{N}} = \text{vol}(N).$$

□

REFERENCES

- [ASS] H. Araki, M-S.B. Smith and L. Smith *On the homotopical significance of the type of von Neumann algebra factors* Commun. Math. Phys. vol 22 (1971) 71-88
- [BFKM] D. Burghelea, L. Friedlander, T. Kappeler and P. McDonald, *Analytic and Reidemeister torsion for representations in finite type Hilbert modules*, to appear in GAFA.
- [BFK] D. Burghelea, L. Friedlander and T. Kappeler, *Torsion for manifolds with boundary and glueing formulae*, preprint 1996.
- [CM] A. Carey and V. Mathai, L^2 Torsion Invariants, Journal of Functional Analysis, **110** (1992) 377-409.
- [CFM] A. Carey, M. Farber and V. Mathai, *Determinant Lines, Von Neumann Algebras and L^2 torsion*, Crelle Journal. **484** (1997) 153-181.
- [Di] J. Dixmier *Von Neumann algebras*, North-Holland, Amsterdam (1981)
- [Do] J. Dodziuk *De Rham-Hodge theory for L^2 cohomology of infinite coverings* Topology **16** (1977) 157-165
- [DM] J. Dodziuk and V. Mathai, *Approximating L^2 invariants for amenable covering spaces: A heat kernel approach*, to appear in "Proceedings of the Bers Colloquium", Contemporary Math.
- [DM2] J. Dodziuk and V. Mathai *Approximating L^2 invariants for amenable covering spaces: A combinatorial approach*, preprint 1996, submitted (see dg-ga archives).
- [FJ] F.T. Farrell and L.E. Jones, *Isomorphism conjectures in algebraic K-theory*, JAMS **6** (1993) 249-298.
- [FK] B. Fuglede and R.V. Kadison, *Determinant theory in finite factors*, Annals of Math. **55** (1952) 520-530
- [L] J. Lott, *Heat kernels on covering spaces and topological invariants*, J. Diff. Geom. **35** (1992) 471-510.
- [Lu] W. Lück, *Approximating L^2 invariants by their finite dimensional analogues*, Geom. and Func. Analysis **4** (1994) 455-481
- [Lu1] W. Lück, *L^2 -Torsion and 3-manifolds*, Conference proceedings and lecture notes in Geometry and Topology, volume III, "Low dimensional Topology", Knoxville, editor: K. Johansson, International Press (1994) 75-107.
- [LuR] W. Lück and M. Rothenberg, *Reidemeister torsion and the K-theory of von Neumann algebras*, K-Theory **5** (1991) 213-264.
- [M] V. Mathai, L^2 analytic torsion, J. Func. Anal. **107** (1992) 369-386
- [Mi] J. Milnor, *Whitehead torsion*, Bull. Amer. Math. Soc., **72**, (1966) 358-426.
- [RS] D. Ray and I. M. Singer, *R-torsion and the Laplacian on Riemannian manifolds*, Advances in Math. **7** (1971) 145-210

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ADELAIDE, ADELAIDE 5005, AUSTRALIA
E-mail address: vmathai@maths.adelaide.edu.au

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, U.S.A.
E-mail address: mel@math.uchicago.edu